

Bald Eagle Wintering Activity Rocky Reach Reservoir



2012 - 2013

**Public Utility District No. 1 of Chelan County
Fish & Wildlife Department
Wenatchee, WA 98807-1231**

INTRODUCTION

The Public Utility District No. 1 of Chelan County (Chelan PUD) owns and operates the Rocky Reach Hydroelectric Project along the Columbia River in central Washington. This project operates under Federal Energy Regulatory Commission (FERC) license #2145-060. Rocky Reach Dam was licensed in 1957 and began generating electricity in 1961. The operating license was renewed in 2009. As prescribed in the Rocky Reach Wildlife Habitat Management Plan required by Article 403 of the Rocky Reach license, Chelan PUD continues to monitor numbers of wintering eagles along Rocky Reach Reservoir (Chelan PUD 2010). This annual report summarizes data collected on wintering bald eagles for the winter of 2012 - 2013.

HISTORY

From 1975 – 1984, the United States Fish and Wildlife Service (USFWS) conducted monthly aerial surveys for wintering bald eagles along the mid-Columbia River, including Rocky Reach Reservoir (Fielder and Starkey 1980). In 1985, Chelan PUD began regularly monitoring eagles during the winter months along Rocky Reach Reservoir.

The bald eagle was delisted on 9 August 2007. As a result, Chelan PUD reduced the monitoring effort from weekly to monthly during the winter months (November – March). This frequency of monitoring exceeds the level suggested in the bald eagle post-delisting monitoring plan (USFWS 2009) and should provide sufficient trend information into the future. Winter surveys allow Chelan PUD to determine distribution of wintering eagles along the Reservoir and to determine the age ratio of sub-adults: adults.

STUDY AREA

Rocky Reach Dam is located approximately 7 miles north of the city of Wenatchee, Washington along the Columbia River at river mile 473.5. The pool behind the dam (Reservoir) extends 42 river miles to Wells Dam (operated by Douglas County PUD) at river mile 515.5. Chelan and Douglas counties border the west and east shores of the reservoir, respectively.

The Reservoir is considered run-of-river with relatively limited water storage capability and active river currents, particularly in the deep main channel of the Columbia River. The Reservoir also has a combination of shallow bays, islands and island complexes, parks, residential, agricultural and natural habitats. Where natural habitat exists, steep cobble or dirt banks comprise much of the reservoir shoreline. Shrub steppe vegetation, fruit orchards, and residential or industrial areas occupy areas up-slope from the riparian edge of the river. Shrub steppe habitat of central Washington is dominated by big sagebrush (*Artemisia tridentata*), rabbitbrush (*Chrysothamnus* spp.), and bluebunch wheatgrass (*Pseudoroegneria spicata*) and is interspersed with pines such as ponderosa pines (*Pinus ponderosa*) and Douglas-fir (*Pseudotsuga menziesii*), especially along the hills to the west that comprise the Eastern Cascades Physiographic province (Franklin and Dyrness 1973). Cottonwoods (*Populus balsamifera* ssp. *trichocarpa*) of differing age groups can also be found along the shores of the river.

SURVEY METHODS

Since golden eagles are a sensitive species in Washington State, both bald and golden eagles are documented during winter surveys. Eagle surveys were conducted by boat monthly from November 2012 – March 2013. A minimum of 2 observers were used on each survey. Surveyors on each side (port and starboard) of the boat monitored the respective shoreline (i.e., Chelan or

Douglas) and both shorelines were surveyed concurrently while traveling upriver along Rocky Reach Reservoir.

Eagles are identified to species (bald or golden) and age class whenever possible. Bald eagles were classified as an adult if they exhibited an all-white head or as a sub-adult in any other color phase. Golden eagles were also aged based on plumage characteristics as described in Wheeler & Clark (1995). If age or species of eagles could not be identified, they were recorded as unknown species or unclassified age status. The location of eagles observed was recorded to the 1/10th river mile as well as bank (Chelan or Douglas) and the perch substrate.

Monthly eagle surveys begin at the Lincoln Rock State Park boat launch and proceeded along the Douglas County shoreline downriver to Rocky Reach Dam, then across the Reservoir to the Chelan County side and upriver along the Chelan County Shore to the upriver end of Turtle Rock Island (river mile 476). At this point the route crosses back over the main channel and downriver along the west side of Turtle Rock Island. At the downriver end of Turtle Rock Island the boat traveled back upriver, along the east shoreline of Turtle Rock Island and the Douglas County shoreline. From the upriver end of Turtle Rock Island, the survey route then continued upriver to the tailrace at Wells Dam. The river just below Wells Dam is quite wide, making it difficult for surveyors to see adequately from the middle of the river. Therefore, the boat cruised near the Douglas County shore beginning at river mile 515.0. Surveyors focused solely on the Douglas County shoreline at this point, continuing upriver to a safe distance below Wells Dam. After the tailrace area was thoroughly surveyed, the Chelan County shoreline was surveyed downriver to river mile 514.9 where the survey then terminated.

RESULTS

During the 2012 - 2013 winter period (November – March), a total of 182 eagles were observed along Rocky Reach Reservoir. Of these, 154 were bald eagles, 26 were golden eagles, and 2 eagles were unidentified. Peak numbers of eagles were observed during January 2013, when a total of 73 eagles (62 bald, 10 golden, and 1 unknown) were observed (Table 1).

Age Ratios

Knowing age ratios of wintering bald eagle populations is important because wintering areas with high proportions of sub-adult bald eagles often indicate an easily available food source which improves the survival chances of younger eagles (Stalmaster 1980, Fitzner et al. 1980, Fielder 1982, Bennetts and McClelland 1991). The maximum number of sub-adult bald eagles (34) observed was on 14 January 2013. The minimum number of sub-adults (2) was observed on 30 November 2012. The average age ratio of bald eagles wintering along the Reservoir for all surveys was 0.9 sub-adults per adult. However, during 3 surveys, this ratio was equal to or greater than 1.0 (Table 2). Bald eagle observations by age class are represented in Figure 1.

Perch Sites and Distribution

Perch substrate was recorded for all eagles observed. Eagles were most commonly observed perched in cottonwoods, flying, or perched in ponderosa pine during surveys (Table 3). Eagles were distributed fairly evenly along much of the Reservoir. On average, eagles were more frequently observed in the area upstream of Turtle Rock Island between Tenas George Canyon and the Ribbon Cliff vicinity, and also in the Howard Flats vicinity (Corral Creek to Wells Dam) (Figure 2).

National Mid-Winter Bald Eagle Surveys

The January 14, 2013 survey was conducted during the National Mid-Winter Bald Eagle Survey timeframe. Data for the survey was submitted via mail to Robert C. Kuntz II, Washington State coordinator for the national counts. A total of 62 bald eagles, 10 golden eagles, and 1 unidentified eagle were observed during this survey. Chelan PUD has regularly participated in the mid-winter bald eagle counting effort.

DISCUSSION

The average number of all eagles observed per survey during the winter of 2012 - 2013 was 36.4. This is higher than the long-term average (1983 - 2012) of 20.8 eagles per survey. The average number of bald eagles observed per survey was 30.8. The ratio of sub-adult bald eagles observed for the 2012 - 2013 winter period was higher than the long-term average of 0.56 sub-adults per adult. The total number of wintering bald eagles (n = 154) observed during the 2012 - 2013 winter is slightly higher than the long-term annual average (AVG = 151) observed since surveys began in 1983 - 1984 (Figure 3). The maximum number of bald eagles observed during any survey during the winter of 2012 - 2013 (MAX = 62) was well above the average maximum number (AVG MAX = 39) observed during any survey to date. The 14 January 2013 survey yielded the highest number of eagle observations ever recorded during any one survey along Rocky Reach Reservoir.

LITERATURE CITED

- Bennetts, R. E., and B. R. McClelland. 1991. Differences in the distribution of adult and immature bald eagles at an autumn concentration in Montana. *Northwest Sci.* 65:223-230.
- Chelan PUD 2010. Rocky Reach Wildlife Habitat Management Plan. Order approving wildlife habitat management plan pursuant to article 403 approved by the FERC on September 22, 2010 (FERC Issuance 20100922-3034, Docket No. P-2145-107).
- Fielder, P. C. 1982. Food habits of bald eagles along the mid-Columbia River, Washington. *Murrelet* 63:46-50.
- Fielder, P. C. and R. Starkey. 1980. Wintering bald eagle use along the upper Columbia River, Washington. pp 177 - 193 in R. Knight, G. Allen, M. Stalmaster, and C. Servheen, eds. *Proc. Washington Bald Eagle Symp., The Nature Conservancy, Seattle.*
- Fitzner, R. E., W. C. Hanson, D. G. Watson, and W. Rickard. 1980. Bald eagles of the Hanford National Environmental Research Park. Pages 207-218 in R. L. Knight, G. T. Allen, M. V. Stalmaster, and C. W. Servheen, eds. *Proc. Washington bald eagle symposium, 1980. The Nature Conservancy, Seattle, WA.*
- Franklin, J. F. and C. T. Dyrness 1973. Natural vegetation of Oregon and Washington. Gen. Tech. Rep. PNW-GTR-008. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station.
- Stalmaster, M. V. 1980. Management strategies for wintering bald eagles in the Pacific Northwest. Pages 49-67 in R. L. Knight, G. T. Allen, M. V. Stalmaster, and C. W. Servheen, eds. *Proc. Washington bald eagle symposium, 1980. The Nature Conservancy, Seattle, WA.*
- U.S. Fish and Wildlife Service. 2009. Post-delisting Monitoring Plan for the Bald Eagle (*Haliaeetus leucocephalus*) in the Contiguous 48 States. U.S. Fish and Wildlife Service, Divisions of Endangered Species and Migratory Birds and State Programs, Midwest Regional Office, Twin Cities, Minnesota.
- Wheeler, B.K. and W.S. Clark. 1995. A photographic guide to North American raptors. Princeton University Press, Princeton, New Jersey, USA.

TABLES AND FIGURES

Table 1. Rocky Reach Reservoir bald eagle surveys, winter of 2012 - 2013.

Date	Bald eagles			Golden eagles				Unknown	
	Adult	Subad	Total	Adult	Subad	Unkn	Total	Eagle	All Total
30-Nov-12	14	2	16	0	0	0	0	0	16
18-Dec-12	7	4	11	3	0	0	3	1	15
14-Jan-13	28	34	62	10	0	0	10	1	73
26-Feb-13	20	19	39	6	1	0	7	0	46
21-Mar-13	10	16	26	5	1	0	6	0	32
Total	79	75	154	24	2	0	26	2	182
AVG	15.8	15	30.8	4.8	-	-	5.2	0.4	36.4

Table 2. Age ratios of sub-adult to adult bald eagles during the survey period.
Golden and unknown eagles and were excluded from analysis.

Date	Adult	Subad	Total	subadults per adult
30-Nov-12	14	2	16	0.1
18-Dec-12	7	4	11	0.6
14-Jan-13	28	34	62	1.2
26-Feb-13	20	19	39	1.0
21-Mar-13	10	16	26	1.6
Total	79	75	154	0.9
AVG	15.8	15	30.8	0.9

Table 3. Perch detail for all eagles observed along Rocky Reach Reservoir, 2012 - 2013.

Perch detail	Total
Cottonwood	57
Flying	45
Ponderosa pine	26
Siberian elm	21
Rock	15
Utility pole	6
Douglas-fir	5
Lombardy poplar	2
Alder	1
Fence post	1
Nest	1
Stump	1
Willow	1
Grand Total	182

Figure 1. Age classifications of bald eagles along Rocky Reach Reservoir.
Golden and unknown eagles and were excluded from analysis.




Figure 2. Average number of all eagle observations by river mile along Rocky Reach Reservoir.




Figure 3. Cumulative observations of bald eagles per survey season. The long-term average is 151 eagles/winter.
Golden and unknown eagles and were excluded from analysis.

