

**From:** [Cordell-Stine, Kelly](#)  
**To:** [Andrew Fielding](#); [Bitterman, Deborah](#); [Brigitte Ranne](#); [Erik Ellis](#); [Jack Oelfke](#); [James Blanchard](#); [Jesse Mccarty](#); [K Vradenburg](#); [Ken Finicle](#); [Matthew Kerec](#); [Matt Monda](#); [molly\\_boyter@blm.gov](mailto:molly_boyter@blm.gov); [Osborn, Jeff](#); [Patrick Tonasket](#); [Patrick Verhey](#); [Paul Willard](#); [Pope, Von](#); [Roger Christophersen \(roger\\_christophersen@nps.gov\)](mailto:roger_christophersen@nps.gov); [Ron Fox \(ron.fox@dfw.wa.gov\)](mailto:Ron.Fox@dfw.wa.gov); [Smith, Michelle](#); [Sokolowski, Rosana](#); [Steve Lewis](#); [Tracy Hames](#); [Volsen, David P \(DFW\)](#)  
**Subject:** Chelan PUD Bald Eagle Wintering Activity Report  
**Date:** Wednesday, January 27, 2016 9:05:56 AM  
**Attachments:** [RRWF BAEA wintering 15-16.pdf](#)

---

Rocky Reach Wildlife Forum Members,

Please find attached the 2015-16 Bald Eagle Wintering Activity Report for Rocky Reach Reservoir. This report is due to RRWF members by April 15 annually.

Surveys were planned for Nov – Dec of 2015 as outlined in the 2010 – 2015 RRWHMP and for January of 2016, as outlined in the 2016 – 2020 RRWHMP.

For the 2016 – 2020 RRWHMP, surveys are reduced to an annual effort. These annual winter bald eagle counts will be scheduled to coincide with the national mid-winter bald eagle counts.

Please let myself or Von Pope know if you have any questions.

Thank you.

[Kelly A. Cordell](#)  
Wildlife Biologist  
Chelan County PUD  
wk. 509.661.4762

cell 509.669.0694

# **Bald Eagle Wintering Activity Rocky Reach Reservoir**



**2015 - 2016**

**Kelly A. Cordell and Von R. Pope**  
**Public Utility District No. 1 of Chelan County**  
**Fish & Wildlife Department**  
**Wenatchee, WA 98807-1231**

## **INTRODUCTION**

The Public Utility District No. 1 of Chelan County (Chelan PUD) owns and operates the Rocky Reach Hydroelectric Project along the Columbia River in central Washington. This project operates under Federal Energy Regulatory Commission (FERC) license #2145-060. Rocky Reach Dam was licensed in 1957 and began generating electricity in 1961. The operating license was renewed in 2009. As prescribed in the Rocky Reach Wildlife Habitat Management Plan required by Article 403 of the Rocky Reach license, Chelan PUD continues to monitor numbers of wintering eagles along Rocky Reach Reservoir (Chelan PUD 2010). This annual report summarizes data collected on wintering bald eagles for the winter of 2015-16.

## **HISTORY**

From 1975 – 1984, the United States Fish and Wildlife Service (USFWS) conducted monthly aerial surveys for wintering bald eagles along the mid-Columbia River, including Rocky Reach Reservoir (Fielder and Starkey 1980). In 1988, Chelan PUD began regularly monitoring eagles during the winter months along Rocky Reach Reservoir.

The bald eagle was delisted from the Endangered Species Act (ESA) on 9 August 2007. As a result, Chelan PUD reduced the monitoring effort from bi-weekly to monthly during the winter months (November – March) beginning in the winter of 2007-08. The frequency of monitoring exceeded the level suggested in the bald eagle post-delisting monitoring plan (USFWS 2009) and provided sufficient trend information. Winter surveys allow Chelan PUD to determine distribution of wintering eagles along the Reservoir and to determine the age ratio of sub-adults: adults. For the 2010 – 2015 Rocky Reach Wildlife Habitat Management Plan (RRWHMP), surveys were conducted monthly. With the recent approval of the 2016 – 2020 RRWHMP, surveys are to be conducted once per winter during January to coincide with the National Mid-winter Bald Eagle Survey (Chelan PUD 2015a).

In 2008, following the delisting of bald eagles from the Endangered Species Act, Chelan PUD reduced the frequency of winter bald eagle monitoring from bi-weekly to monthly. In 2010, the first Rocky Reach Wildlife Habitat Management Plan was approved and directed Chelan PUD to continue winter bald eagle monitoring as conducted during the first license term.

In 2015, the second Rocky Reach Wildlife Habitat Management Plan was submitted for FERC approval. The plan was approved on December 1, 2015, and reduced the number of winter bald eagle surveys from monthly to one time each winter, beginning in 2016. Annual winter bald eagle counts will be scheduled to coincide with the national mid-winter bald eagle counts.

This report summarizes the monitoring effort conducted under the first Rocky Reach Wildlife Habitat Management Plan for the winter of 2015-2016.

## **STUDY AREA**

Rocky Reach Dam is located approximately 7 miles north of the city of Wenatchee, Washington along the Columbia River at river mile 473.5. The pool behind the dam (Reservoir) extends 42 river miles north to Wells Dam (operated by Douglas County PUD) at river mile 515.5. Chelan and Douglas counties border the west and east shores of the reservoir, respectively.

The Reservoir is considered run-of-river with relatively limited water storage capability and active river currents, particularly in the deep main channel of the Columbia River. The Reservoir also has a combination of shallow bays, islands and island complexes, parks, residential, agricultural and natural habitats. Where natural habitat exists, steep cobble or dirt banks comprise much of the reservoir shoreline. Shrub steppe vegetation, fruit orchards, and residential or industrial areas occupy areas up-slope from the riparian edge of the river. Shrub steppe habitat of central Washington is dominated by big sagebrush (*Artemisia tridentata*), rabbitbrush (*Chrysothamnus* spp.), and bluebunch wheatgrass (*Pseudoroegneria spicata*) and is interspersed with pines such as ponderosa pines (*Pinus ponderosa*) and Douglas-fir (*Pseudotsuga menziesii*), especially along the hills to the west that comprise the Eastern Cascades Physiographic province (Franklin and Dyrness 1973). Cottonwoods (*Populus balsamifera* ssp. *trichocarpa*) of differing age groups can also be found along the shores of the river.

## **SURVEY METHODS**

Since golden eagles are a sensitive species in Washington State, both bald and golden eagles are documented during winter surveys. Eagle surveys were to be conducted by boat monthly from November and December 2015, and once during January 2016 to coincide with the dates for the national Mid Winter Bald Eagle Survey. A minimum of 2 observers were used on each survey. Surveyors on each side (port and starboard) of the boat monitored the respective shoreline (i.e., Chelan or Douglas) and both shorelines were surveyed concurrently while traveling upriver along Rocky Reach Reservoir.

Eagles are identified to species (bald or golden) and age class (adult or sub-adult) whenever possible. Bald eagles were classified as an adult if they exhibited an all-white head or as a sub-adult in any other color phase. Golden eagles were also aged based on plumage characteristics as described in Wheeler & Clark (1995). If age or species of eagles could not be identified, they were recorded as unknown species and/or unclassified age. The location of eagles observed was recorded to the 1/10<sup>th</sup> river mile as well as bank (Chelan or Douglas) and the perch substrate.

Monthly eagle surveys begin at the Lincoln Rock State Park boat launch and proceeded along the Douglas County shoreline downriver to Rocky Reach Dam, then across the Reservoir to the Chelan County side and upriver along the Chelan County Shore to the upriver end of Turtle Rock Island (river mile 476). At this point the route crosses back over the main channel and downriver along the west side of Turtle Rock Island. At the downriver end of Turtle Rock Island the boat traveled back upriver, along the east shoreline of Turtle Rock Island and the Douglas County shoreline. From the upriver end of Turtle Rock Island, the survey route then continued upriver to the tailrace at Wells Dam. The river just below Wells Dam is quite wide, making it difficult for surveyors to see adequately from the middle of the river. Therefore, the boat cruised near the Douglas County shore beginning at river mile 515.0. Surveyors focused solely on the Douglas County shoreline at this point, continuing upriver to a safe distance below Wells Dam. After the tailrace area was thoroughly surveyed, the Chelan County shoreline was surveyed downriver to river mile 514.9 where the survey then terminated.

## **RESULTS**

During the 2015-16 winter period (November – January), a total of 2 surveys were completed. The November survey was canceled due to poor weather conditions. A total of 87 eagles were observed along Rocky Reach Reservoir throughout the survey season. Of these, 77 were bald eagles, 9 were

golden eagles, and 1 eagle was unidentified. Peak numbers of eagles were observed during January 2016, when a total of 48 eagles (48 bald and 0 golden) were observed (Table 1).

### **Age Ratios**

Knowing age ratios of wintering bald eagle populations is important because wintering areas with high proportions of sub-adult bald eagles often indicate an easily available food source which improves the survival chances of younger eagles (Stalmaster 1980, Fitzner et al. 1980, Fielder 1982, Bennetts and McClelland 1991). The maximum number of sub-adult bald eagles (24) observed occurred during the January survey. The minimum number of sub-adults (13) occurred during the December survey. The average age ratio of bald eagles wintering along the Reservoir for both surveys was 0.9 sub-adults per adult (Table 2). Bald eagle observations by age class are represented in Figure 1.

### **Perch Sites and Distribution**

Perch substrate was recorded for all eagles observed. Eagles were most commonly observed perched in cottonwoods, on cliffs or rocky outcrops, or ponderosa pines during surveys (Table 3). Eagles were distributed fairly evenly along much of the Reservoir. On average, eagles were more frequently observed in the area between Spencer Canyon and Knapp Tunnel, and also in the Howard Flats vicinity (Corral Creek to Wells Dam) (Figure 2).

### **National Mid-Winter Bald Eagle Surveys**

The 8 January 2016 survey was conducted during the National Mid-Winter Bald Eagle Survey timeframe (preferred target dates 8 – 9 January 2016). The National Mid-Winter Bald Eagle survey is overseen by the United States Geological Survey (USGS) and in coordination with the United States Army Corps of Engineers (ACOE). Data for the survey was submitted via email on 12 January 2016 to Stephen Lesky, ACOE, Washington State coordinator for the national counts. A total of 48 bald eagles and no golden eagles were observed during this survey. Chelan PUD has regularly participated in the mid-winter bald eagle counting effort.

### **Long-term Summary**

Over a period of 28 years, (winter of 1988-89 to winter of 2015-16) the average numbers of bald eagles observed wintering along Rocky Reach Reservoir has increased from an average of 16.1 bald eagles/survey (1988-89) to 38.5 bald eagles per survey (2015-16). This increasing trend is shown in Figure 4. The average number of bald eagles observed over this 28 year period is 19.5 bald eagles per survey. The maximum number of wintering bald eagles is typically observed during the month of January or February, with much lower maximum numbers in November, December, and March (Figure 5). The maximum count by month for wintering bald eagles has occurred during January in 12 years (42.8%), during February in 11 years (39.3%), and the same high count was observed during both months in 5 years (17.9%). No maximum counts occurred during November, December or March from 1988-89 through 2014-15. For the 28 year period, the month of January has the highest average number of wintering bald eagles observed with 32.9 bald eagles per survey followed closely by February with 31.8 bald eagles per survey.

## **DISCUSSION**

The average number of all eagles (bald, golden, and unidentified) observed per survey during the winter of 2015-16 was 43.5. This is higher than the long-term average (1988 - 2015) of 19.4 eagles per survey. The average number of bald eagles observed per survey during 2015-16 was 38.5. The

average number of golden eagles observed per survey was 4.5. The ratio of sub-adult: adult bald eagles observed for the 2015-16 winter period ( $n = 0.9$ ) was higher than the long-term average of 0.56 sub-adults per adult. The total number of wintering bald eagles ( $n = 77$ ) observed during the 2015-16 winter is lower than the long-term annual average (AVG = 149) observed since surveys began in 1988-89 (Figure 3). However, fewer surveys ( $n = 2$ ) were conducted during the 2015-16 season as a result of the reduction in survey effort under the 2016 – 2020 RRWHMP. The maximum number of all eagles observed during any survey during the winter of 2015-16 (MAX = 48) was slightly above the average maximum number (AVG MAX = 40) observed during any survey to date.

In consultation with the Rocky Reach Wildlife Forum, Chelan PUD reduced its wintering bald eagle effort in 2008 following the delisting of the bald eagle from the ESA in 2007. This change was to continue winter monitoring with reduced effort during the 5-year period following delisting of the bald eagle. Survey effort during the winter months (November – March) was reduced from bi-weekly to monthly during the 2010 – 2015 RRWHMP (Chelan PUD 2010). As the 5-year period following delisting has passed and wintering bald eagle numbers continue to increase, along with increasing numbers of nesting bald eagles along Rocky Reach Reservoir (Chelan PUD 2015b), the RRWF reduced the number of wintering surveys from once monthly to once per winter for the 2016 – 2020 RRWHMP. The single winter count shall coincide with the National Mid-Winter Bald Eagle survey. Chelan PUD has submitted January survey data annually since 2008 for Rocky Reach Reservoir in conjunction with the National Mid-Winter survey effort. The January effort will ensure the peak of the winter season is surveyed and the data set will continue to be shared with the National Mid-Winter Bald Eagle survey coordinators.

## LITERATURE CITED

- Bennetts, R. E., and B. R. McClelland. 1991. Differences in the distribution of adult and immature bald eagles at an autumn concentration in Montana. *Northwest Sci.* 65:223-230.
- Chelan PUD 2015a. Rocky Reach Wildlife Habitat Management Plan 2016 – 2020. License Article 403. Rocky Reach Hydroelectric Project. Commission Project No. 2145. September 15, 2015. Public Utility District No. 1 of Chelan County. Wenatchee, WA 98801.
- Chelan PUD 2015b. Bald eagle nesting activity. Rocky Reach Reservoir. Public Utility District No. 1 of Chelan County. Wenatchee, WA 98801.
- Chelan PUD 2010. Rocky Reach Wildlife Habitat Management Plan. Order approving wildlife habitat management plan pursuant to article 403 approved by the FERC on September 22, 2010 (FERC Issuance 20100922-3034, Docket No. P-2145-107).
- Fielder, P. C. 1982. Food habits of bald eagles along the mid-Columbia River, Washington. *Murrelet* 63:46-50.
- Fielder, P. C. and R. Starkey. 1980. Wintering bald eagle use along the upper Columbia River, Washington. pp 177 - 193 in R. Knight, G. Allen, M. Stalmaster, and C. Servheen, eds. *Proc. Washington Bald Eagle Symp., The Nature Conservancy, Seattle.*
- Fitzner, R. E., W. C. Hanson, D. G. Watson, and W. Rickard. 1980. Bald eagles of the Hanford National Environmental Research Park. Pages 207-218 in R. L. Knight, G. T. Allen, M. V. Stalmaster, and C. W. Servheen, eds. *Proc. Washington bald eagle symposium, 1980. The Nature Conservancy, Seattle, WA.*
- Franklin, J. F. and C. T. Dyrness 1973. Natural vegetation of Oregon and Washington. Gen. Tech. Rep. PNW-GTR-008. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station.
- Stalmaster, M. V. 1980. Management strategies for wintering bald eagles in the Pacific Northwest. Pages 49-67 in R. L. Knight, G. T. Allen, M. V. Stalmaster, and C. W. Servheen, eds. *Proc. Washington bald eagle symposium, 1980. The Nature Conservancy, Seattle, WA.*
- U.S. Fish and Wildlife Service. 2009. Post-delisting Monitoring Plan for the Bald Eagle (*Haliaeetus leucocephalus*) in the Contiguous 48 States. U.S. Fish and Wildlife Service, Divisions of Endangered Species and Migratory Birds and State Programs, Midwest Regional Office, Twin Cities, Minnesota.
- Wheeler, B.K. and W.S. Clark. 1995. A photographic guide to North American raptors. Princeton University Press, Princeton, New Jersey, USA.

**TABLES AND FIGURES**

**Table 1. Rocky Reach Reservoir bald eagle surveys, winter of 2015 – 2016.**

<b>Date</b>	<b>Bald eagles</b>			<b>Golden eagles</b>				<b>Unknown</b>			
	<b>Adult</b>	<b>Subad</b>	<b>Total</b>	<b>Adult</b>	<b>Subad</b>	<b>Unkn</b>	<b>Total</b>	<b>Eagle</b>	<b>All Total</b>	<b>subs/adult</b>	<b>adults/sub</b>
No November Survey											
22-Dec-15	16	13	29	7	2	0	9	1	39	0.8	1.2
8-Jan-16	24	24	48	0	0	0	0	0	48	1.0	1.0
<b>Total</b>	<b>40</b>	<b>37</b>	<b>77</b>	<b>7</b>	<b>2</b>	<b>0</b>	<b>9</b>	<b>1</b>	<b>87</b>	<b>0.9</b>	<b>1.1</b>
<b>AVG</b>	<b>20.0</b>	<b>18.5</b>	<b>38.5</b>	<b>3.5</b>	<b>0.0</b>	<b>0.0</b>	<b>4.5</b>	<b>0.5</b>	<b>43.5</b>	<b>0.9</b>	<b>1.1</b>



**Table 2. Age ratios of sub-adult to adult bald eagles during the winter of 2015 – 2016.**  
Golden and unknown eagles and were excluded from analysis.

<b>Date</b>	<b>Adult</b>	<b>Sub-adult</b>	<b>Total</b>	<b>Sub-adults per adult</b>
<i>No November Survey</i>				
22-Dec-15	16	13	29	0.8
8-Jan-16	24	24	48	1.0
<b>Total</b>	<b>40</b>	<b>37</b>	<b>77</b>	<b>0.9</b>
<b>AVG</b>	<b>20</b>	<b>18.5</b>	<b>38.5</b>	<b>0.9</b>

**Table 3. Perch detail for bald eagles observed along Rocky Reach Reservoir during 2015 – 2016.**

<b>Perch detail</b>	<b>Total</b>
Cottonwood	24
Flying (not perched)	14
Rock	11
Ponderosa pine	10
Siberian elm	7
Douglas-fir	5
Utility pole	3
Alder	2
Lombardy poplar	1
<b>Total</b>	<b>77</b>

**Figure 1. Age classification of bald eagles along Rocky Reach Reservoir during the winter of 2015 – 16.**

Golden and unknown eagles and were excluded from analysis.




Figure 2. Average number of eagle observations (all species) by river mile along Rocky Reach Reservoir during the winter of 2015 – 16.




Figure 3. Maximum count of bald eagles obtained during the month of January, 1988 – 2015. The average maximum/survey during the January effort is 32 bald eagles. Golden and unknown eagles and were excluded from analysis.

### January Max Count




Figure 4. The average number of wintering bald eagles observed along Rocky Reach Reservoir from 1988 – 2015 and the long term linear average showing the increase over the 28 year period.




Figure 5. Average maximum number of bald eagles by month along Rocky Reach Reservoir, 1988 – 2015.

