

**A Monitoring Study to Quantify Dam Passage and Tributary  
Escapement of Adult Pacific Lamprey in the Rocky Reach Project  
Area and the Mid-Columbia River**

**SUBMITTED TO:**

**Steve Hemstrom  
Public Utility District No.1 of Chelan County  
327 North Wenatchee Avenue  
Wenatchee, WA 98807**

**SUBMITTED BY:**

**Julie Maenhout  
Blue Leaf Environmental  
2301 West Dolarway Road, Suite 3  
Ellensburg, Washington, 98926**

**PRINCIPAL PROPONENT:**

**Mark Timko, Blue Leaf Environmental**

**October 4, 2017**



**BLUE LEAF  
ENVIRONMENTAL**

## Executive Summary

On February 19, 2009, Public Utility District No. 1 of Chelan County (Chelan PUD) filed application for a new Federal Energy Regulatory Commission (FERC) operating license for the Rocky Reach Hydroelectric Project (Project), which included a Pacific Lamprey Management Plan (PLMP) (FERC 2009). On May 1, 2009, FERC issued an “order modifying and approving” the PLMP.

The goal of the PLMP is to achieve No Net Impact (NNI) on Pacific lamprey by measuring ongoing Project-related impacts, if any, then implementing appropriate and reasonable measures to reduce or eliminate effects of those Project operations on lamprey, and finally, implementing on-site or off-site measures to address unavoidable effects. The Chelan PUD PLMP is intended to be consistent with other management plans in the mid-Columbia region.

The goals and objectives of the PLMP are to be achieved through Protection, Mitigation, and Enhancement (PME) measures when unavoidable Project effects are measured. In support of these PME’s, this passage monitoring study was designed to address PME’s 4.1, measure any Project-related impacts on upstream and downstream passage of adult Pacific lamprey and eliminate those impacts to the extent appropriate and reasonable; 4.1.5, evaluation of upstream passage modification; and 4.1.6, determination of any significant adult downstream passage effects.

This study used adult lamprey collected at Priest Rapids Dam (Grant PUD) and implanted with FDX-PIT tags to address the following objectives for Chelan PUD’s study:

1. Continue to assess adult Pacific lamprey passage rates and behavior at Rocky Reach Dam using FDX- PIT technology to evaluate the in-fishway passage improvements Chelan PUD completed for lamprey in 2011.
2. Estimate adult Pacific lamprey passage success at Rocky Reach Dam to allow comparison to passage rates at other Projects on the Columbia River, per the PLMP.
3. Estimate the proportion of volitional escapement into mid-Columbia River tributaries.
4. Determine final return location, if possible, for the proportion of fish detected in the Wenatchee River, the Entiat River, at mid-Columbia dams upstream of Rocky Reach Dam, and for fish that are detected at Rocky Reach Dam but ultimately do not pass upstream through the dam’s fishway.
5. Estimate the proportion of fish that overwinter in the mainstem Columbia River based on following-year detections of fish at mainstem dams and any following-year first detections in tributaries.
6. Assist and coordinate with Douglas PUD to provide Douglas PUD with up to 50 fish captured for Chelan PUD’s study to enable Douglas to double tag lamprey (FDX-PIT tag and active acoustic tag) to assess movement through Rocky Reach Reservoir (Chelan PUD objective) and approach behavior up to and through Wells Dam tailrace and Wells Dam (Douglas PUD objective). These fish will be released upstream of Rocky Reach Dam but downstream of the confluence of the Entiat River to allow volitional escapement of lamprey into the Entiat River.

Two hundred eleven adult Pacific lampreys were collected from Priest Rapids Dam fishways during the 2016 peak migration period, implanted with full-duplex PIT tags, and released above Rock Island Dam at Kirby Billingsley Hydro Park (RM 461.8) in five release days between 3 August and 17 August. The fish were distributed approximately 90% near the right (west) bank and 10% along the left (east) bank to approximate Rock Island Dam fishway use and top of ladder exit proportions. Existing PIT tag detection infrastructure at Rocky Reach Dam, and other mainstem Columbia River dams and in tributary streams was used to monitor the migration behavior of tagged individuals in the study area. The PIT Tag Information System (PTAGIS) was queried to find detection records for tagged individuals. A detection history was created for each tagged lamprey and used to calculate metrics related to the study objectives.

Of the 211 lampreys released, 174 (82.5%) have been detected in either the adult fishway at Rocky Reach or Wells dams or in a mid-Columbia River tributary, 19 (90.5%) from the left bank releases and 155 (81.6%) from the right bank releases. As of September 18, 2017, the Rocky Reach fishway passage rate was 98.8%. One hundred sixty-two of the 164 fish detected in the Rocky Reach fishway were last detected at the fishway exit and two were last detected at the lower weir antennas in the upper fishway. Of the fish that passed Rocky Reach dam in 2016, one of these fish passed Wells Dam and was last detected in the Chewuch River above Winthrop, WA in 2016, another fish was last detected at Wells Dam in 2017, 13 fish were subsequently detected entering the Entiat River in 2017, five were last detected at the Rocky Reach Dam Juvenile Fish Facility in 2017, while the remaining 143 were last detected at the Rocky Reach fishway exit in 2016. Eleven fish were last detected in the Wenatchee River, four in 2016 and seven in 2017. No lampreys from the 2016 study have been detected at Rock Island Dam. The median travel rate to reach the exit of the Rocky Reach fishway for fish released on the left bank was 5.8 rkm/day (range 1.2-23.8 rkm/day) compared to 3.4 rkm/day (range 0.4-23.3 rkm/day) for right bank releases. While the median travel rate from release to first detection in the Wenatchee River was 26.6 rkm/day (range 1.1-28.9 rkm/day). The single fish that was last detected upstream of Wells Dam in the Chewuch River exhibited an overall travel rate of 5.6 rkm/day from release to first detection in the Chewuch River. We note that travel times for lamprey moving into tributaries is based on small sample sizes. Freshwater maturity time is also unknown. Therefore, at this point we caution the comparison of lamprey travel rates. Travel rates were not calculated for 2017 detections because it is unknown when those fish resumed their migration after overwintering. Of the 211 lampreys released, 87.7% were either last detected at Rocky Reach Dam or not detected at all after release, indicating possible mainstem spawning in the Rock Island or Rocky Reach reservoirs. It is also possible that lamprey may have either moved upstream past Wells Dam, downstream through Rock Island Dam, or into the Wenatchee or Entiat rivers undetected.

1.0	INTRODUCTION .....	6
1.1	Adult Lamprey Dam Passage.....	6
2.0	STUDY METHODS .....	8
2.1	Collection.....	8
2.2	Tagging .....	9
2.3	Releases.....	10
2.4	Monitoring .....	11
2.5	Analysis of Detections .....	15
3.0	RESULTS .....	15
4.0	DISCUSSION .....	18
5.0	REFERENCES .....	19

**LIST OF FIGURES**

Figure 1.	Holding containers (coolers) plumbed with flow-through river water at the Priest Rapids Dam left bank holding facility.....	9
Figure 2.	Girth was measured at the anterior insertion of the first dorsal fin. Length, inter-dorsal distance (IDD) to the nearest mm, and weight in grams, was measured and recorded before each lamprey was tagged.....	10
Figure 3.	PIT tagged adult Pacific lamprey release sites in Rock Island Reservoir at Kirby Billingsley Hydro Park for 2016 Rocky Reach lamprey passage study. 10% (n=21) of all tagged lamprey for the study were released near the left bank and 90% (n=190) on the right bank as shown. These proportions emulate proportions of lamprey using the left bank and right bank fishways at Rock Island Dam.....	11
Figure 4.	Full-duplex PIT tag detection sites are located at Rock Island, Rocky Reach, and Wells dams as well as tributary detection sites in the Wenatchee and Entiat rivers. Detections of tagged lamprey released for the study were queried using the PTAGIS website ( <a href="http://www.ptagis.org">www.ptagis.org</a> ).....	12
Figure 5.	Detail of full-duplex PIT tag detection sites located at Rocky Reach (top) and Wells (bottom) dams. Detections of tagged lamprey released for the study were queried using the PTAGIS website ( <a href="http://www.ptagis.org">www.ptagis.org</a> ). .....	13
Figure 6.	Detail of full-duplex PIT tag detection sites located at the lower Wenatchee (top) and Entiat (bottom) rivers. Detections of tagged lamprey released for the study were queried using the PTAGIS website ( <a href="http://www.ptagis.org">www.ptagis.org</a> ).....	14

Figure 7. Number of tagged lamprey that passed Rocky Reach Dam per week in 2016. The dots represent when the two fish that did not pass were last detected in the upper fishway. No additional fish were detected passing Rocky Reach Dam in 2017. .... 16

LIST OF TABLES

Table 1. The total quantity of lamprey tagged and released by release date in August 2016. .... 16

Table 2. The last detection sites as of September 18, 2017 for 2016 and 2017 detected lampreys totaling 174 of 211 unique tagged fish. .... 17

Table 3. Travel time, in days, from release sites to the last detection at the exit of the Rocky Reach Dam fishway. .... 17

Table 4. Travel rates, in river kilometers per day, from release sites to the last detection at the exit of the Rocky Reach Dam fishway. .... 17

Table 5. Travel time, in days, from release site to the first detection in the Rocky Reach Dam fishway.. 18

Table 6. Travel time, in hours, from first detection to last detection in the Rocky Reach Dam fishway. . 18

## 1.0 INTRODUCTION

On February 19, 2009, Public Utility District No. 1 of Chelan County (Chelan PUD) filed application for a new Federal Energy Regulatory Commission (FERC) operating license for the Rocky Reach Hydroelectric Project (Project), which included a Pacific Lamprey Management Plan (PLMP) (FERC 2009). On May 1, 2009, FERC issued an “order modifying and approving” the PLMP.

The goal of the PLMP is to achieve No Net Impact (NNI) on Pacific lamprey by measuring ongoing Project-related impacts, if any, then implementing appropriate and reasonable measures to reduce or eliminate effects of those Project operations on lamprey, and finally, implementing on-site or off-site measures to address unavoidable effects. The Chelan PUD PLMP is intended to be consistent with other management plans in the mid-Columbia region.

The goals and objectives of the PLMP are to be achieved through a series of Protection, Mitigation, and Enhancement (PME) measures. In support of these PME’s, this passage monitoring study was designed to address PME’s 4.1, measure any Project-related impacts on upstream and downstream passage of adult Pacific lamprey and eliminate those impacts to the extent appropriate and reasonable; 4.1.5, evaluation of upstream passage modification; and 4.1.6, determination of any significant adult downstream passage effects.

### 1.1 Adult Lamprey Dam Passage

Pacific Lamprey (*Entosphenus tridentatus*) numbers have declined since the 1940’s as a result of many factors including spawning habitat loss, impediments to migration, changing ocean conditions, and decreased water quality (Close et al. 1995, 2002), and other potentially unrecognized conditions. Much focus over the past 10 years has been placed on studying migrating adults. Modifications have been made to many dam operations and fishway structures to improve lamprey passage. Passive tags, such as half-duplex passive integrated transponder tags (HDX- PIT), and active tags, such as radio tags (RT) or acoustic tags (i.e. Juvenile Salmonid Acoustic Transmitter System [JSATS]) have been employed to monitor adult lamprey passage primarily at dams in the lower Columbia River (Keefer et al. 2009a, Keefer et al. 2009b, Keefer et al. 2009c, Noyes et al. 2014). In 2015, a ban implemented in 2004 on the use of full-duplex PIT tags (FDX-PIT) for lamprey research was lifted by the PIT tag steering committee (PTAGIS 2015). This change made the existing FDX-PIT tag detection infrastructure common in mainstem Columbia and Snake river dams and tributaries available to lamprey researchers. One of the advantages of using FDX-PIT tags for lamprey research is the use of the PIT Tag Information System (PTAGIS) for querying detections that may occur throughout monitored sites in the Columbia and Snake river drainages. This makes it possible to imply escapement rates into tributaries or “final fate” of lamprey that are detected in spawning tributaries. Deciphering final return locations has been more difficult to quantify using other passive tag technologies.

Areas within upstream fishways can be problematic for lamprey passage (Moser et al. 2003) and as such, Chelan PUD initiated a study in 2004 using radio-telemetry to assess lamprey passage behavior and success through the Rocky Reach Dam upstream fishway. The estimated net ladder passage efficiency (NLPE), or proportion of fish that were detected in the tailrace of the dam that exited the upstream fishway was 47%, which was comparable to studies at other Columbia River dams at the time (Stevenson et al. 2005). In 2011, Chelan PUD performed

significant modifications to the Rocky Reach fishway intended to aid lamprey passage. In 2014, Chelan PUD completed a study to assess passage behavior of adult lamprey using HDX-PIT tags and found that net ladder passage, or the proportion of fish with final successful passage was estimated at 66% (Blue Leaf Environmental 2015) though the most upstream HDX antenna system in Rocky Reach fishway was compromised during a critical two-week period in the study. Although the 2004 and 2014 studies employed different tag technologies and reported slightly differing passage metrics, the studies suggest that lamprey passage through Rocky Reach Dam has improved, perhaps significantly, since the completion of in-fishway passage improvements in 2011. Due to limited detection capabilities of HDX-PIT tags in tributaries, it is unknown what proportion of fish released for the 2014 study ascended spawning tributaries. Review of total year fishway counts of lamprey passing Rock Island Dam and Rocky Reach Dam suggests that dam to dam passage conversion rates since 2011 are high for Rocky Reach (Chelan PUD 2015 unpublished data).

## **1.2 Study objectives**

The study described herein used adult lamprey collected at Priest Rapids Dam and implanted with FDX-PIT tags to address the following objectives:

1. Assess adult Pacific lamprey passage rates and behavior at Rocky Reach Dam using FDX- PIT technology to evaluate the in-fishway passage improvements Chelan PUD completed for lamprey in 2011.
2. Estimate adult Pacific lamprey passage rate success at Rocky Reach Dam to allow comparison to passage rates at other Projects on the Columbia River, per the PLMP.
3. Estimate the proportion of volitional escapement into mid-Columbia River tributaries.
4. Determine final return locations, if possible, for the proportion of fish detected in the Wenatchee River, the Entiat River, at mid-Columbia dams upstream of Rocky Reach Dam, and for fish that are detected at Rocky Reach Dam but ultimately do not pass upstream through the dam's fishway.
5. Estimate the proportion of fish that overwinter in the mainstem Columbia River based on following-year detections of fish at mainstem dams and any following-year first detections in tributaries.
6. Assist and coordinate with Douglas PUD to provide Douglas PUD with up to 50 fish captured for Chelan PUD's study to enable Douglas to double tag lamprey (FDX-PIT tag and active acoustic tag) to assess movement through Rocky Reach Reservoir (Chelan PUD objective) and approach behavior up to and through Wells Dam tailrace and Wells Dam (Douglas PUD objective). These fish will be released upstream of Rocky Reach Dam but downstream of the confluence of the Entiat River to allow volitional escapement of lamprey into the Entiat River.

### **1.3 Study Area**

The study area is defined by the FERC licensed project area which includes Rocky Reach Dam and the associated tailrace area and reservoir. The project area is referenced here as the Rocky Reach Project (RRP). The Rocky Reach Reservoir spans approximately 66.6 km (41.4 miles) of the Columbia River. The Rocky Reach Reservoir has a full-pool (707.0 ft msl) surface area of approximately 8,828 acres. The RRP has only one major tributary input, the Entiat River, but both the Wenatchee River and the Entiat River were included in the study area because FDX-PIT monitoring arrays exist in both, and estimation of tributary escapement rates was desired. Both rivers were monitored for lamprey detections during the study.

## **2.0 STUDY METHODS**

### **2.1 Collection**

In 2016, adult Pacific lamprey were collected using mechanical fish traps located in the lower fishways at Priest Rapids Dam from 2 August to 17 August. All fish were scanned for an existing PIT tag. Previously tagged fish were released upstream of Priest Rapids Dam. All other fish were transported to a holding facility at Priest Rapids Dam left bank. Fish were placed in 20.5 gallon insulated containers (coolers) plumbed with flow-through river water, with a maximum of 15 fish per container (Figure 1).





Figure 1. Holding containers (coolers) plumbed with flow-through river water at the Priest Rapids Dam left bank holding facility.

## 2.2 Tagging

All lamprey were tagged with a FDX-PIT tag (12.5 mm L x 2.1 mm Dia, 0.1 g). At the beginning of each tagging and release day, the appropriate number of tags were placed in a Nolvasan (Chlorhexidine diacetate) solution for ten minutes then rinsed with distilled water. Injection needles were also disinfected in a Nolvasan solution for ten minutes and then held in trays filled with distilled water. Injection needles were disinfected and rinsed in the same manner between each fish. Physical metrics and anesthetic times for each fish were recorded on data sheets.

One lamprey at a time was transferred from a container by hand, using water soaked cotton gloves, to a heavy anesthetic bath with 120 parts per million (ppm) clove oil, until the fish lost equilibrium and swimming motion ceased (approximately 5 minutes). Start and end time (i.e., anesthetic time) was recorded. Once the lamprey was fully anesthetized, length, girth, inter-dorsal distance (IDD) to the nearest mm, and weight in grams, was measured and recorded

(Figure 2). A FDX-PIT tag was scanned into an ACCESS database. The FDX-PIT tag was then injected using a Biomark MK10 implanter and N125 injector needle above the ventral midline, in the softer, thinner part of the abdomen between the ventral midline and the muscular “rib cage”. The injection site was in line with the anterior insertion of the first dorsal fin. The lamprey was then scanned to confirm PIT tag ID, and transferred to a holding/transport container on flow-through river water, labeled with the appropriate release location and number, and container number. There were no more than 12 lampreys per release container. Lampreys were held in the holding/transport containers until fully recovered from anesthetic, and released later the same day.



Figure 2. Girth was measured at the anterior insertion of the first dorsal fin. Length, inter-dorsal distance (IDD) to the nearest mm, and weight in grams, was measured and recorded before each lamprey was tagged.

### 2.3 Releases

After recovery, fish were transported in aerated coolers filled with river water to Kirby Billingsley Hydro Park (RM 461.8) and transferred to the custody of Chelan PUD fisheries staff.

Chelan PUD staff transferred the transport coolers to a boat. The fish were released into the Columbia River, distributed approximately 90% near the right (west) bank and 10% along the left (east) bank to approximate the number of lamprey using each of the Rock Island Dam fishways and top of ladder exit proportions based on total lamprey counts (Figure 3).



Figure 3. PIT tagged adult Pacific lamprey release sites in Rock Island Reservoir at Kirby Billingsley Hydro Park for 2016 Rocky Reach lamprey passage study. 10% (n=21) of all tagged lamprey for the study were released near the left bank and 90% (n=190) on the right bank as shown. These proportions emulate proportions of lamprey using the left bank and right bank fishways at Rock Island Dam.

## 2.4 Monitoring

This study employed passive FDX-PIT tags and existing FDX-PIT detection arrays installed at fixed locations (Rocky Reach Dam adult fishway, Wells Dam adult fishways, and multiple locations in the Wenatchee and Entiat rivers), to determine passage success and tributary escapement of adult Pacific lamprey in the RRP (Figure 4). Tagged fish were monitored as they moved past fixed FDX-PIT tag detection arrays located in the adult fishway at Rocky Reach and Wells dams and also in mid-Columbia River tributaries such as the Wenatchee, Entiat, and Methow rivers (Figure 5 and Figure 6). Detections histories of study fish were assembled by querying the PTAGIS database at appropriate intervals. Queries were used to look for detections in the expected locations as well as system-wide in the event that study fish did not behave as expected.




Figure 4. Full-duplex PIT tag detection sites are located at Rock Island, Rocky Reach, and Wells dams as well as tributary detection sites in the Wenatchee and Entiat rivers. Detections of tagged lamprey released for the study were queried using the PTAGIS website ([www.ptagis.org](http://www.ptagis.org)).



### Rocky Reach Dam Adult Fishway (RRF)

PIT Tag Antenna Map: PTAGIS Cnfg. #100; March 2006



PIT Tag Information Systems  
Columbia Basin

### Wells Dam: Wells Fish Ladders (WEA)

Interrogation Coil Map Revised: February, 2002 v.1.0  
Orifice Dimensions: 21" wide x 34.5" high




Figure 5. Detail of full-duplex PIT tag detection sites located at Rocky Reach (top) and Wells (bottom) dams. Detections of tagged lamprey released for the study were queried using the PTAGIS website ([www.ptagis.org](http://www.ptagis.org)).





Figure 6. Detail of full-duplex PIT tag detection sites located at the lower Wenatchee (top) and Entiat (bottom) rivers. Detections of tagged lamprey released for the study were queried using the PTAGIS website ([www.ptagis.org](http://www.ptagis.org)).

## 2.5 Analysis of Detections

Detection records from the PTAGIS database were compiled in a relational database maintained by Blue Leaf Environmental (BLE) staff. A detection history was generated for each tag. Detection records were used to determine behavioral characteristics of tagged lamprey in the study area. Key metrics included the proportion of tagged fish that entered the Wenatchee or Entiat rivers based on instream detections, the proportion of tagged fish that passed through the Rocky Reach Dam adult fishway, and the proportion detected at or in the Wells Dam adult fishway, and in the Methow Basin. Also, the movement rate from release to detection at the above locations was estimated. Additionally, we report the proportion of fish that overwintered in the study area and resumed movements in 2017.

## 3.0 RESULTS

A total of 211 adult lampreys were tagged with FDX-PIT tags in five tag-days, and released in the same days between 3 August and 17 August (Table 1). Of the 211 lampreys released, 174 (82.5%) have been detected in either the adult fishway at Rocky Reach Dam or Wells Dam, or in a mid-Columbia River tributary. In total, 19 lampreys were detected (90.5%) from the left bank releases and 155 (81.6%) from the right bank releases.

As of September 18, 2017, the Rocky Reach fishway passage rate at study completion is 98.8%. One hundred sixty-two of the 164 fish detected in the Rocky Reach fishway were last detected at the fishway exit and two were last detected at the lower weir antennas in the upper fishway (antenna A3 and A4; Figure 5). Fish began passing Rocky Reach Dam within one day of being released at Kirby Billingsley Hydro Park and a majority (94.4%) of fish passed within a two-week period in the month of August (Figure 7). No additional fish were detected passing Rocky Reach Dam in 2017 that were not detected in 2016. Of the fish that passed Rocky Reach Dam, one of these fish passed Wells Dam and was last detected in the Chewuch River above Winthrop, WA in 2016, another fish was last detected in the Wells Dam fishway in 2017, 13 fish were subsequently detected entering the Entiat River in 2017, and five fish were last detected at the Rocky Reach Dam Juvenile Fish Bypass Surface Collector in 2017. Eleven fish were last detected in the Wenatchee River, four in 2016 and seven in 2017 (Table 2). The remaining 143 fish were last detected at the Rocky Reach fishway exit in 2016. One of these fish passed Rocky Reach Dam in August of 2016 but then was last detected in the Wenatchee River a full year later in July, 2017. The confluence of the Wenatchee and Columbia rivers, at RM 468.5, is 7.5 RM upstream from the release site and 4.5 RM downstream of Rocky Reach Dam.

Table 1. The total quantity of lamprey tagged and released by release date in August 2016.

Release Date	Left Bank	Right Bank	Total
8/3/2016	5	41	46
8/5/2016	6	52	58
8/10/2016	3	36	39
8/12/2016	4	33	37
8/17/2016	3	28	31
Total	21	190	211




Figure 7. Number of tagged lamprey that passed Rocky Reach Dam per week in 2016. The dots represent when the two fish that did not pass were last detected in the upper fishway. No additional fish were detected passing Rocky Reach Dam in 2017.



Table 2. The last detection sites as of September 18, 2017 for 2016 and 2017 detected lampreys totaling 174 of 211 unique tagged fish.

Last Detection Site	2016	2017	Total
Wenatchee River	4	7	11
Rocky Reach Dam Adult Fishway	143	0	143
Rocky Reach Dam JFB Surface Collector	0	5	5
Entiat River	0	13	13
Wells Dam Adult Ladder	0	1	1
Chewuch River (Methow)	0	1	1
Total	147	27	174

Travel times from release to passage through Rocky Reach Dam (release to last detection at the Rocky Reach fishway exit) varied by release site. Median travel times were 3.3 days for left bank released fish (n=19) and 5.7 days for right bank released fish (n=143) (Table 3). The median travel rate from release to the exit of the Rocky Reach fishway for left bank releases was 5.8 rkm/day (range 1.2-23.8 rkm/day) compared to 3.4 rkm/day (range 0.4-23.3 rkm/day) for right bank releases (Table 4). The smaller sample size of left bank released fish may account for the reduced variation in travel times and rates compared to the right bank released fish. The single fish that was last detected upstream of Wells Dam in the Chewuch River exhibited a travel time of 18.5 days and travel rate of 4.8 rkm/day from release to first detection in the Wells Dam fishway, with an overall travel time of 32.6 days and travel rate of 5.6 rkm/day from release to first detection in the Chewuch River. The median travel time from release to detection in the Wenatchee river was 0.6 days (range 0.5-13.6 days) with a median travel rate of 26.6 rkm/day (range 1.1-28.9 rkm/day). We note that travel times for lamprey moving into tributaries is based on small sample sizes. Freshwater maturity time is also unknown. Therefore, at this point we caution the comparison of lamprey travel rates. Travel rates were not calculated for 2017 detections because it is unknown when those fish resumed their migration after overwintering.

Table 3. Travel time, in days, from release sites to the last detection at the exit of the Rocky Reach Dam fishway.

Travel Time (days)						
Release Site	n	Min	Max	Median	Mean	SD
Left Bank	19	0.8	15.7	3.3	5.1	4.1
Right Bank	143	0.8	43.4	5.7	7.9	7.4

Table 4. Travel rates, in river kilometers per day, from release sites to the last detection at the exit of the Rocky Reach Dam fishway.

Travel Rate (rkm/d)						
Release Site	n	Min	Max	Median	Mean	SD
Left Bank	19	1.2	23.8	5.8	7.2	5.8
Right Bank	143	0.4	23.3	3.4	5.3	4.4

A majority of the total travel time exhibited by tagged lampreys that passed Rocky Reach Dam is accounted for in time from release to first detection within the Rocky Reach fishway, indicating lamprey spent little time in the upper portion of the fishway. The median travel time from release to first detection in the Rocky Reach fishway for fish released on the left bank was 1.8 days (19 fish, range 0.8-15.7 days) compared to 3.7 days (145 fish range 0.8-21.6 days) for right bank releases (Table 5). Travel time from first detection in the fishway to last detection in the fishway is better represented in hours due to the minimum and median travel times being less than one hour. Substantial variation was observed in times of individual fish, but the median travel time was 0.4 and 0.6 hours for left and right bank releases, respectively (Table 6).

Table 5. Travel time, in days, from release site to the first detection in the Rocky Reach Dam fishway.

Travel Time (days)						
Release Site	n	Min	Max	Median	Mean	SD
Left Bank	19	0.8	15.7	1.8	4.4	4.1
Right Bank	145	0.8	21.6	3.7	5.6	4.9

Table 6. Travel time, in hours, from first detection to last detection in the Rocky Reach Dam fishway.

Travel Time (hours)						
Release Site	n	Min	Max	Median	Mean	SD
Left Bank	19	0.1	141.4	0.4	17.3	38.4
Right Bank	145	0.1	882.5	0.6	56.9	134.4

#### 4.0 DISCUSSION

Some studies have reported that lamprey rarely travel long distances after overwintering to spawn (Courter et al. 2012; McIlraith et al. 2015), while others report lamprey moving significant distances downstream to spawn after overwintering (Starcevich et al. 2014). There is also some indication that lamprey may engage in mainstem spawning, particularly in the tailrace of hydropower dams where spawning habitat known to be used by fall Chinook salmon occurs (Dauble et al. 1999; McMichael et al. 2005). Overall, a large percentage of the lamprey in this study may have engaged in mainstem spawning within the Rocky Reach Reservoir. Only one PIT-tagged fish which entered the Chewuck River is known to have passed Wells Dam.

As of September 2017, 87.7% of the tagged lampreys were either last detected at Rocky Reach Dam or not detected at all after release. The five lamprey last detected at the Rocky Reach Juvenile Fish Bypass Surface Collector in late June and early July of 2017 were most likely post-spawn and drifting downstream. Although, based on the early summer detection times, it's possible they were traveling downstream after overwintering, in search of a place to spawn. It is also possible that some lamprey may have moved upstream to Wells Dam, downstream through Rock Island Dam, or into the Wenatchee or Entiat rivers undetected. A majority of the

movement into tributaries occurred in the spring of 2017 when flows were well above the 10-year average (<http://www.cbr.washington.edu/dart/dart/html>). High flow is known to cause decreased detection efficiency at in-river PIT tag interrogation systems (Connolly et al. 2008).

Passage rates at Rocky Reach Dam reported in this study are among the highest reported on the mainstem Columbia River to date, and are a significant increase from the 2004 radio telemetry estimate and 2014 half-duplex PIT tag estimate (Stevenson et al. 2005; BLE 2015). Of the lamprey detected in the Rocky Reach fishway, 98.8% were last detected at the exit in the 2016 migration year. Interestingly, passage estimates at the lower Columbia River dams were also among the highest estimated in previous studies as well, at 60%, 58%, and 73% for Bonneville, The Dalles, and John Day dams, respectively (Keefer et al. 2015).

Lamprey travel times from release to the exit of the Rocky Reach fishway in this study were similar to the HDX study conducted in 2014. The median travel time for 2014 HDX tagged fish was 1.7 days compared to 1.8 days and 3.7 days for left and right bank released fish in this study, respectively (BLE 2015; Table 5).

## 5.0 REFERENCES

- Blue Leaf Environmental. 2015. HDX PIT results 2014-2015 at Rocky Reach Dam. Technical memorandum prepared for Chelan Public Utility District, Wenatchee, WA. Prepared by R. R. O'Connor Nov 13, 2015.
- Chelan PUD. 2015. Unpublished data. Pacific lamprey fishway window counts and unadjusted conversion rates between Rock Island Dam and Rocky Reach Dam, 2011-2015.
- Close, D. A., M. S. Fitzpatrick and H. W. Li. 2002. The ecological and cultural importance of a species at risk of extinction, Pacific lamprey. *Fisheries* 27:19-25.
- Close, D.A., M.S. Fitzpatrick, H.W. Li., B.L. Parker, D.R. Hatch and G.A. James. 1995. Status report of the Pacific Lamprey (*Lampetra tridentata*) in the Columbia River Basin. Bonneville Power Administration, Portland, Oregon.
- Connolly, P.J., I.G. Jezorek, K. Martens, and E.F. Prentice. 2008. Measuring performance of two stationary interrogation systems for detecting downstream and upstream movement of PIT-tagged salmonids. *North American Journal of Fisheries Management* 28:402-417.
- Courter, I., S. Duery, J. Vaughan, C. Peery, M. Morasch, R. McCoun, B. Clemens, and C. Schreck. 2012. Migration behavior and distribution of adult Pacific lamprey in the Willamette Basin. Prepared for The Columbia River Inter-Tribal Fish Commission, Portland, Oregon.
- Dauble, D. D., R. L. Johnson, and A. P. Garcia. 1999. Fall Chinook salmon spawning in the tailraces of lower Snake River hydroelectric projects. *Transactions of the American Fisheries Society* 128:672-679.
- Keefer, M. L., C. T. Boggs, C. A. Peery, M. L. Moser. 2009a. Adult Pacific lamprey migration in the lower Columbia River: 2007 radiotelemetry and half-duplex PIT tag studies. Technical Report 2009-1 of Idaho Cooperative Fish and Wildlife Research Unit to U.S. Army Corps of Engineers, Portland District.

- Keefer, M. L., M. L. Moser, C. T. Boggs, W. R. Daigle, and C. A. Peery. 2009b. Variability in migration timing of adult Pacific lamprey (*Lampetra tridentata*) in the Columbia River, U.S.A. *Environmental Biology of Fishes* 85:253-264.
- Keefer, M. L., M. L. Moser, C. T. Boggs, W. R. Daigle, and C. A. Peery. 2009c. Effects of body size and river environment on the upstream migration of adult Pacific lamprey. *North American Journal of Fisheries Management* 29:1214-1224.
- Keefer, M. L., C. C. Caudill, E. L. Johnson, T. S. Clabough, M. A. Jepson, C. J. Noyes, C. T. Boggs, M. A. Kirk, S. C. Corbett, K. E. Frick, and M. L. Moser. 2015. Adult Pacific lamprey migration in the Columbia and Snake rivers: 2014 radiotelemetry and half-duplex PIT tag studies and retrospective summaries. Draft Technical Report 2015-12 of University of Idaho to U.S. Army Corps of Engineers, Portland District.
- McIlraith, B. J., C. C. Caudill, B. P. Kennedy, C. A. Peery, and M. L. Keefer. 2015. Seasonal migration behaviors and distribution of adult Pacific lampreys in unimpounded reaches of the Snake River Basin. *North American Journal of Fisheries Management* 35:123-134.
- McMichael, G. A., C. A. McKinstry, J. A. Vucelick, and J. A. Lukas. 2005. Fall Chinook salmon spawning activity versus daylight and flow in the tailrace of a large hydroelectric dam. *North American Journal of Fisheries Management* 25:573-580.
- Moser, M.L., R.W. Zabel, B.J. Burke, L.C. Stuehrenberg, and T.C. Bjornn. 2003. Factors affecting adult Pacific lamprey passage rates at hydropower dams: using “time to event” analysis of radiotelemetry data. In Spedicato, M.T., G. Lembo, and G. Marmulla (editors) *Aquatic telemetry: advances and applications*. Proceedings of the Fifth Conference on Fish Telemetry held in Europe. Ustica, Italy, 9-13 June 2003.
- Noyes, C.J., C.C. Caudill, T.S. Clabough, D.C. Joosten, and M.L. Keefer. 2014. Adult Pacific lamprey migration behavior and escapement in the Bonneville Reservoir and lower Columbia River monitored using the Juvenile Salmonid Acoustic Telemetry System (JSATS), 2013. Technical Report #2014-7. Prepared for the U.S. Army Corps of Engineers Portland District, Portland, OR.
- PTAGIS. 2015. PIT Tag Information System newsletter. February 2015, Volume 13, Issue 1. Pp. 3-4.
- Starcevich, S.J., S.L. Gunckel, and S.E. Jacobs. 2014. Movements, habitat use, and population characteristics of adult Pacific lamprey in a coastal river. *Environmental Biology of Fishes* 97:939-953.
- Stevenson, J.R., P. Westhagen, D.J. Snyder, J.R. Skalski, and A.E. Giorgi. 2005. Evaluation of adult Pacific lamprey passage at Rocky Reach Dam using radiotelemetry techniques, 2004. Technical Report prepared by BioAnalysts, Inc. for Chelan Public Utility District, Wenatchee, WA.